

It's about the moments

Continue enjoying moments with loved ones, while treating your chronic hepatitis B



Not actual patients.

VEMLIDY is a prescription medicine used to treat chronic (long-lasting) hepatitis B virus (HBV) in adults with stable (compensated) liver disease.

**Talk to your doctor today
about starting VEMLIDY**

 **Vemlidy**[®]
tenofovir alafenamide 25mg
tablets

[Click here](#) for Important Facts about VEMLIDY, with **important warnings on worsening of hepatitis B infection.**

Every moment is precious with your family and friends.

That's why it's important to treat your chronic (long-lasting) hepatitis B virus, or HBV. VEMLIDY is a prescription medicine that treats chronic hepatitis B in adults with stable (compensated) liver disease. VEMLIDY may lower the amount of HBV in your body and improve the condition of your liver.

What is the most important information I should know about VEMLIDY?

VEMLIDY can cause serious side effects, including:

- **Worsening of hepatitis B virus (HBV) infection.** Your HBV infection may get worse (flare-up) if you take VEMLIDY and then stop taking it. A “flare-up” is when your HBV infection suddenly returns in a worse way than before. Do not stop taking VEMLIDY without first talking to your healthcare provider, as they will need to monitor your health regularly to check your liver.

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.



What does the liver do?

The liver is one of the most important parts of the body. It does a lot of things that help you every day, including:



Fighting infection



Helping digest food



Storing energy and nutrients



Removing harmful substances from the blood



Not an actual patient.

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.

 **Vemlidy**[®]
tenofovir alafenamide 25mg tablets

What is chronic hepatitis B?

Chronic hepatitis B is a disease caused by long-term infection by the hepatitis B virus. Over time, it can lead to serious liver damage.

It's important to know that you **can't** get hepatitis B from touching, kissing, drinking alcohol, food, water, air, or sharing cups and utensils

What are the symptoms?

Hepatitis B symptoms may include:



Tiredness/
fatigue



Fever



Joint pain



Stomach pain



Nausea and
vomiting



Loss of appetite



Yellow eyes
or skin
(known as jaundice)



Light-colored
stools
or dark urine



Many patients with chronic hepatitis B do not experience any symptoms. **Even if you don't feel any symptoms, the hepatitis B virus is still in your body and may be harming your liver**

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.

 **Vemlidy**[®]
tenofovir alafenamide 25mg tablets

How chronic hepatitis B can damage the liver

The higher your viral load, the higher the risk for you to develop liver damage.

Chronic hepatitis B can lead to:



Inflammation

Swelling of the liver that can cause scarring over time



Fibrosis

Scarring of the liver where scar tissue replaces healthy tissue



Cirrhosis

Severe scarring that may affect how the liver works



Liver Cancer

The longer you have hepatitis B, the more damage it can cause. Because of the risk for severe liver damage, it is important to see your doctor regularly, even if you feel fine

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.

 **Vemlidy**[®]
tenofovir alafenamide 25mg tablets



Not actual patients.

The goal of HBV treatment

The goal of prescription medicine is to:



Decrease the amount of virus in the body



Reduce the risk of liver damage

If you are currently on treatment, there are tests that can tell how well your medicine is working. Talk to your doctor about why these tests are important and how to schedule them.

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.

 **Vemlidy**[®]
tenofovir alafenamide 25mg tablets

Experience each moment




Not actual patients.

What is VEMLIDY?

VEMLIDY is a prescription medicine used to treat chronic (long-lasting) hepatitis B virus (HBV) in adults with stable (compensated) liver disease.

- VEMLIDY may lower the amount of HBV in your body.
- VEMLIDY may improve the condition of your liver.

Timeline of FDA Approvals: Oral Antiviral Treatments for Chronic Hep B



This timeline is not intended to compare the efficacy or safety of the products listed above.

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.

 **Vemlidy**®
tenofovir alafenamide 25mg tablets

Studies in a broad range of patients have shown that VEMLIDY:



reduces the amount of hepatitis B in your body (viral load)



may improve the condition of the liver

What should I tell my healthcare provider before taking VEMLIDY?

All of your medical conditions, including if you have end stage renal disease (ESRD) or HIV-1 infection. Your healthcare provider may test you for HIV infection before starting VEMLIDY. If you have HIV and only take VEMLIDY, the HIV virus may develop resistance and become harder to treat.

If you are pregnant or plan to become pregnant. It is not known if VEMLIDY will harm your unborn baby. Tell your healthcare provider if you become pregnant during treatment with VEMLIDY.

If you are breastfeeding or plan to breastfeed. It is not known if VEMLIDY passes into your breast milk. Talk with your healthcare provider about the best way to feed your baby.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines may affect how VEMLIDY works. Keep a list of all your medicines and show it to your healthcare provider and pharmacist. **Do not start a new medicine without telling your healthcare provider.** Ask your healthcare provider if it is safe to take VEMLIDY with all of your other medicines.

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.

 **Vemlidy**[®]
tenofovir alafenamide 25mg tablets

Same dose every day with VEMLIDY



Pill shown is actual size.
(8 mm in diameter)

VEMLIDY is one small tablet that you take once a day with food.

You must take VEMLIDY exactly how your doctor tells you to. If you are on dialysis, on your dialysis days, take VEMLIDY after dialysis. It's also important to take VEMLIDY on a regular dosing schedule to avoid missing doses.

Do not stop taking VEMLIDY unless your doctor tells you to. If you stop taking your medicine, your HBV infection could get worse (flare-up). A "flare-up" is when your HBV infection suddenly returns in a worse way than before.

It is very important that you refill your prescription on time, before you run out. When your VEMLIDY supply starts to run low, get more from your doctor or pharmacy. This will help make sure that you are not at risk for missing any doses and causing a flare-up.

If you take too much VEMLIDY, call your healthcare provider or go to the nearest hospital emergency room right away.

Tips to help you remember



Make it a habit by taking your medicine at the same time every day



Set an alarm or download a reminder app on your phone



Get help from your family or friends

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.

 **Vemlidy**[®]
tenofovir alafenamide 25mg tablets



Not an actual patient.

What are the possible side effects of VEMLIDY?

- **New or worse kidney problems, including kidney failure.** Your healthcare provider may do blood and urine tests to check your kidneys when starting and during treatment with VEMLIDY. Your healthcare provider may tell you to stop taking VEMLIDY if you develop new or worse kidney problems.
- **Too much lactic acid in your blood (lactic acidosis),** which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feeling dizzy or lightheaded, or a fast or abnormal heartbeat.
- **Severe liver problems,** which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark “tea-colored” urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.

The most common side effect of VEMLIDY is headache.

These are not all the possible side effects of VEMLIDY. Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.



Worried about cost?

You may be eligible for the VEMLIDY Co-pay Coupon Program

The VEMLIDY Co-pay Coupon Program is here to help reduce your out-of-pocket costs

- Out-of-pocket costs will be covered up to \$5,000 each year
- No monthly limit
- For eligible patients who have commercial insurance

As low as



Translation services are available in a variety of languages. Please call **1-877-627-0415** today to get started

*See specific terms and conditions at www.vemlidy.com.

Gilead's Advancing Access® Program is here to help you

- Do you need help understanding your health insurance and Gilead medication costs?
- Even if you are not eligible for the Co-pay Coupon Program you can call Advancing Access at the number below to learn about other financial resources that may be available to assist with treatment and insurance costs

For more information, call **1-800-226-2056**

Hours: Monday-Friday, 9 AM to 8 PM, ET



Multilingual help is available. Please notify one of our associates if non-English assistance is needed. We will do our best to accommodate your language needs

Please visit www.vemlidy.com for more information on eligibility and enrollment.

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.



(VEM-lih-dee)

IMPORTANT FACTS

This is only a brief summary of important information about VEMLIDY. This does not replace talking to your healthcare provider about your condition and your treatment.

MOST IMPORTANT INFORMATION ABOUT VEMLIDY

VEMLIDY may cause serious side effects, including:

- **Worsening of hepatitis B virus (HBV) infection.** Your HBV infection may suddenly get worse (flare-up) if you stop taking VEMLIDY. Do not run out of VEMLIDY. Do not stop taking VEMLIDY without first talking to your healthcare provider, as they will need to check your health regularly after stopping therapy.

ABOUT VEMLIDY

VEMLIDY is a prescription medicine used to treat chronic (long-lasting) hepatitis B virus (HBV) in adults with stable (compensated) liver disease.

VEMLIDY:

- May lower the amount of HBV in your body.
- May improve the condition of your liver.

BEFORE TAKING VEMLIDY

Tell your healthcare provider about all of your medical conditions, including if you:

- Have HIV-1 infection. Your healthcare provider may test you for HIV infection before you start using VEMLIDY. If you have both HBV and HIV, and you only take VEMLIDY, the HIV virus may develop resistance and be harder to treat.
- Have end stage renal disease (ESRD).
- Are pregnant or plan to become pregnant.
- Are breastfeeding or plan to breastfeed.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that interact with VEMLIDY.

POSSIBLE SIDE EFFECTS OF VEMLIDY

VEMLIDY can cause serious side effects, including:

- Those in the “Most Important Information About VEMLIDY” section.
- New or worse kidney problems, including kidney failure. Your healthcare provider may do blood and urine tests to check your kidneys when starting and during treatment with VEMLIDY.
- Too much lactic acid in your blood (lactic acidosis), which is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feeling dizzy or lightheaded, or a fast or abnormal heartbeat.
- Severe liver problems, which in rare cases can lead to death. Tell your healthcare provider right away if you get these symptoms: skin or the white part of your eyes turns yellow, dark “tea-colored” urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.

The most common side effect of VEMLIDY is headache.

These are not all the possible side effects of VEMLIDY. Tell your healthcare provider right away if you have any side effect that bothers you or that does not go away.

HOW TO TAKE VEMLIDY

Take VEMLIDY 1 time each day with food.

GET MORE INFORMATION

- This is only a brief summary of important information about VEMLIDY. Talk to your healthcare provider or pharmacist to learn more.
- Go to [VEMLIDY.com](https://vemlidy.com) or call 1-800-GILEAD-5
- If you need help paying for your medicine, visit [VEMLIDY.com](https://vemlidy.com) for program information.



Not an actual patient.

***Speak with your doctor about VEMLIDY
at your next visit***



To learn more about VEMLIDY,
please visit www.vemlidy.com

[Click here](#) for Important Facts about VEMLIDY, including **important warnings**.

VEMLIDY, the VEMLIDY Logo, ADVANCING ACCESS, GILEAD, and the GILEAD Logo are trademarks of Gilead Sciences, Inc., or its related companies. All other trademarks referenced herein are the property of their respective owners.

©2021 Gilead Sciences, Inc. All rights reserved.
VEMC0194 03/21

